

UNIVERSITY OF TORONTO

Faculty of Arts and Science  
Department of Ecology and Evolutionary Biology



Mathematical modeling in ecology and evolution  
(EEB314/1430)

*Test 1 – September 22, 2025*

*Available Time: 50m*

*Permitted Materials: Pen/pencil*

**Question 1** ( 13/13 points ) Transposable elements are small segments of DNA that selfishly replicate themselves within a genome, often at a cost to the individual. They are found across the tree of life to a widely varying degree, e.g., they make up about 85% of the maize (corn) genome but only about 2% of the baker's yeast genome. To better understand the factors determining the number of transposable elements in a genome, Charlesworth and Charlesworth (1983) built a model. The model assumes that the transposable elements replicate at per capita rate  $u$  and are lost at per capita rate  $v$ , giving growth rate  $\lambda = u - v$ , which we will assume is positive. It also assumes there are  $T$  places in a genome where the transposable elements can exist and the fitness of an individual with  $n$  transposable elements in their genome is  $w_n$ . Finally, fitness is assumed to decline linearly with the number of transposable elements,  $w_n = 1 - ns$ , each reducing fitness by amount  $s$ . Putting this all together, they determined that the mean number of transposable elements per individual changed at rate

$$\frac{d\bar{n}}{dt} = \bar{n}\lambda - \bar{n}\left(1 - \frac{\bar{n}}{T}\right)\left(\frac{s}{1 - \bar{n}s}\right).$$

- (a) ( 3 points ) Show how to find the two equilibria of this model,  $\hat{n} = 0$  and

$$\hat{n} = \frac{\lambda - s}{s(\lambda - 1/T)}.$$

We'll refer to these as the zero and nonzero equilibria, respectively.

- (b) ( 3 points ) Show how to determine that the nonzero equilibrium is biologically valid when either

(a)  $1/T \leq s \leq \lambda$  or

(b)  $\lambda \leq s \leq 1/T$ .

- (c) ( 4 points ) Using the product and chain rules we have

$$\begin{aligned} \frac{d}{d\bar{n}} \left( \frac{d\bar{n}}{dt} \right) &= \lambda - \left(1 - \frac{\bar{n}}{T}\right) \left( \frac{s}{1 - \bar{n}s} \right) \\ &\quad - \bar{n} \left( \frac{s}{1 - \bar{n}s} \right) \left( \left(1 - \frac{\bar{n}}{T}\right) \left( \frac{s}{1 - \bar{n}s} \right) - \frac{1}{T} \right). \end{aligned}$$

Show how to determine that stability of the zero equilibrium requires  $\lambda < s$  and stability of a valid nonzero equilibrium requires  $1/T \leq s < \lambda$ . (Hint: from part a you may have noticed that

$$\left(1 - \frac{\bar{n}}{T}\right) \left( \frac{s}{1 - \bar{n}s} \right) = \lambda$$

at the nonzero equilibrium.)

- (d) ( 2 points ) Under this model, what needs to be true for an initially rare transposable element to be able to increase but not increase indefinitely?

- (e) ( 1 points ) Assuming transposable elements are at the nonzero equilibrium, which of the following may explain the larger number of transposable elements in maize vs. baker's yeast?
- (a) transposable elements have a smaller  $s$  in maize than yeast
  - (b) transposable elements have a larger  $T$  in maize than yeast

———— *End of Exam* ————